

## 32-1127: EB13 Recombinant Protein

**Alternative Name :** Interleukin-27 subunit beta,IL-27 subunit beta,IL-27B,Epstein-Barr virus-induced gene 3 protein,EBV-induced gene 3 protein,EB13,IL27B.

### Description

Source : Escherichia Coli. EB13 Human Recombinant produced in E.Coli is a single, non-glycosylated, Polypeptide chain containing 209 amino acids fragment (21-229) having a molecular weight of 23.3kDa. The EB13 is purified by proprietary chromatographic techniques. EB13 has an induced expression in B lymphocytes in reaction to Epstein-Barr virus infection. EB13 encodes a secreted glycoprotein belonging to the hematopoietin receptor family, and heterodimerizes with a 28 kDa protein to form iIL-27. EB13 drives rapid clonal expansion of naive cd4(+) t-cells. EB13 strongly synergizes with IL-12 to activate IFN-gamma production of naive cd4(+) t-cells. EB13 mediates its biologic effects through the cytokine receptor wsx-1/tccr.

### Product Info

**Amount :** 10 µg  
**Purification :** Greater than 90% as determined by(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.  
**Content :** EB13 Human Recombinant was lyophilized from a solution containing 10mM Acetic Acid and 0.5% Mannitol.  
**Storage condition :** Lyophilized EB13 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution EB13 should be stored at 4°C between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.  
**Amino Acid :** RKGPPAALTLPVQCASRYPIAVDCSWTLPPAPNSTSPVSFIATYRLGMAARGHSWPCLQQTPTSTSTCTITDVQLFSMAPYVLNVTAVHPWGSSSFVPFITEHIIKPDPEGVRLSPLAERQLQVQWEPGSPFPFEIFSLKYWIRYKRQGAARFHRVGPPEATSFILRAVRPRARYVQVAAQDLTDYGELSDWSLPATATMSLGK.

### Application Note

It is recommended to reconstitute the lyophilized EB13 in sterile 10mM Acetic acid not less than 100µg/ml, which can then be further diluted to other aqueous solutions. Assay data for Human recombinant EB13 is based upon qualitative binding to anti-EB13 antibody.

