

32-1130: EG VEGF Recombinant Protein

Alternative Name : PK1,PRK1,Prokineticin 1,EG-VEGF.

Description

Source : Escherichia Coli. EG-VEGF Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 86 amino acids and having a molecular mass of 9605 Dalton. The EG-VEGF is purified by proprietary chromatographic techniques. Endocrine gland-derived vascular endothelial growth factor (EG-VEGF) induces proliferation, migration, and fenestration in capillary endothelial cells derived from endocrine glands. Its expression is induced by hypoxia and is restricted to the steroidogenic glands (ovary, testis, adrenal, and placenta). Its expression is often complementary to the expression of VEGF (MIM 192240), suggesting that these molecules function in a coordinated manner. EG-VEGF potently contracts gastrointestinal (gi) smooth muscle. Induces proliferation, migration and fenestration (the formation of membrane discontinuities) in capillary endothelial cells derived from endocrine glands. Has little or no effect on a variety of other endothelial and non-endothelial cell types.

Product Info

Amount :	20 µg
Purification :	Greater than 97.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
Content :	The protein was lyophilized from a concentrated (1mg/ml) solution with no additives.
Storage condition :	Lyophilized EG-VEGF Human Recombinant although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution EG-VEGF should be stored at 4°C between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.
Amino Acid :	AVITGACERD VQCGAGTCCA ISLWLRGLRM CTPLGREGEE CHPGSHKVPF FRKRKHTTCP CLPNLLCSRF PDGRYRCSMD LKNINF.

Application Note

It is recommended to reconstitute the lyophilized Endocrine Gland Vascular Endothelial Growth Factor in sterile 18MΩ[⊙]-cm H₂O not less than 100Ω[⊙]µg/ml, which can then be further diluted to other aqueous solutions.

