

32-1132: EGF Recombinant Protein

Alternative Name : Urogastrone,URG,EGF.

Description

Source : Escherichia Coli. Epidermal Growth Factor Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 53 amino acids and having a molecular mass of 6222 Dalton.The EGF is purified by proprietary chromatographic techniques. Epidermal growth factor has a profound effect on the differentiation of specific cells in vivo and is a potent mitogenic factor for a variety of cultured cells of both ectodermal and mesodermal origin. The EGF precursor is believed to exist as a membrane-bound molecule which is proteolytically cleaved to generate the 53-amino acid peptide hormone that stimulates cells to divide. EGF stimulates the growth of various epidermal and epithelial tissues in vivo and in vitro and of some fibroblasts in cell culture.

Product Info

Amount :	0.5mg
Purification :	Greater than 98.0% as determined by:(a) Analysis by SEC-HPLC.(b) Analysis by SDS-PAGE.
Content :	The protein was lyophilized from a concentrated (1mg/ml) solution containing PBS pH-7.4. Lyophilized Epidermal Growth Factor Recombinant although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution EGF should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
Storage condition :	
Amino Acid :	NSDSECLSH DGYCLHDGVC MYIEALDKYA CNCVVG YIGE RCQYRDLKWW ELR.

Application Note

It is recommended to reconstitute the lyophilized Epidermal Growth Factor in sterile 18M Ω -cm H₂O not less than 100 μ g/ml, which can then be further diluted to other aqueous solutions. The ED₅₀, calculated by the dose-dependant proliferation of murine BALB/c 3T3 cells (measured by 3H-thymidine uptake) is < 0.1 ng/ml corresponding to a specific activity of 1 x 10,000,000 Units/mg.

