

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-1136: mEGF Recombinant Protein

Alternative Name: Urogastrone, URG, EGF.

Description

Source: Escherichia Coli. Epidermal Growth Factor Mouse Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 53 amino acids including 3 intramolecular disulfide-bonds and having a molecular mass of 6 kDa. The EGF is purified by proprietary chromatographic techniques. Epidermal growth factor has a profound effect on the differentiation of specific cells in vivo and is a potent mitogenic factor for a variety of cultured cells of both ectodermal and mesodermal origin. The EGF precursor is believed to exist as a membrane-bound molecule which is proteolytically cleaved to generate the 53-amino acid peptide hormone that stimulates cells to divide. EGF stimulates the growth of various epidermal and epithelial tissues in vivo and in vitro and of some fibroblasts in cell culture.

Product Info

Amount: 0.5mg

Purification: Greater than 98.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Content: The protein was lyophilized with no additives.

 $Lyophilized\ Epidermal\ Growth\ Factor\ Recombinant\ although\ stable\ at\ room\ temperature\ for\ 3$

weeks, should be stored desiccated below -18°C. Upon reconstitution EGF should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended

to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Amino Acid: NSYPGCPSSY DGYCLNGGVC MHIESLDSYT CNCVIGYSGD RCOTRDLRWW ELR.

Application Note

Storage condition:

It is recommended to reconstitute the lyophilized Epidermal Growth Factor in sterile 18MÃ[]Â[]c-cm H2O not less than 100Ã[]Â[]A[]g/ml, which can then be further diluted to other aqueous solutions. The activity is determined by the dose-dependent proliferation of mouse BALB/c 3T3 cells and is typically less than 0.1ng/ml.

