

32-1205: Flt3 Ligand HEK Recombinant Protein

Alternative Name : Fms-related tyrosine kinase 3 ligand,FLK2,STK1,CD135,Stem Cell Tyrosine Kinase 1,FLT3LG,Flt3.

Description

Source : HEK293 (Human Embryonic Kidney cell line). Flt3-Ligand Human Recombinant produced in HEK293 cells is a glycosylated, polypeptide chain, migrates as a diffuse band on SDS-PAGE due to heterogeneous glycosylation at 30kDa. Flt3-Ligand is purified by proprietary chromatographic techniques. FLT3 ligand is a receptor for the fl cytokine has a tyrosine-protein kinase activity & a growth factor that regulates proliferation of early hematopoietic cells. Flt3-Ligand synergizes with other CSFs and interleukins to induce growth and differentiation.

Product Info

| | |
|----------------------------|---|
| Amount : | 2 µg |
| Purification : | "Greater than 95.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE." |
| Content : | The protein was lyophilized with no additives. |
| Storage condition : | Lyophilized Flt3-Ligand although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Flt3-Ligand should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles. |
| Amino Acid : | TQDCSFQHSP ISSDFAVKIR ELSDYLLQDY PVTVASNLQD EELCGGLWRL VLAQRWMERL KTVAGSKMQG LLERVNTEIH FVTKCAFQPP PSCLRFVQTN ISRLLQETSE QLVALKPWIT RQNFSRCLEL QCQPDSSTLP PPWSPRPLEA TAPTAPQP. |

Application Note

It is recommended to reconstitute the lyophilized Flt3-Ligand in sterile 18MΩ-cm H₂O not less than 100µg/ml, which can then be further diluted to other aqueous solutions. The ED₅₀ as determined by the dose dependent proliferation of human OCI-AML5 cells is 0.5 to 1.0 ng/ml (Specific Activity: 2.0 x 1,000,000 – 1.0 x 1,000,000 units/mg).

