

## 32-1209: rmFlt3 Ligand Recombinant Protein

**Alternative Name :** Fms-related tyrosine kinase 3 ligand,FLK2,STK1,CD135,Stem Cell Tyrosine Kinase 1,FLT3LG,Flt3.

### Description

Source : Escherichia Coli. Flt3 Ligand Rhesus Macaque Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 159 amino acids and having a molecular mass of 18.0kDa. The Flt3 Ligand Rhesus Macaque is purified by proprietary chromatographic techniques. FLT3 ligand is a receptor for the fl cytokine has a tyrosine-protein kinase activity & a growth factor that regulates proliferation of early hematopoietic cells. Flt3-Ligand synergizes with other CSFs and interleukins to induce growth and differentiation.

### Product Info

<b>Amount :</b>	10 µg
<b>Purification :</b>	Greater than 97.0% as determined by SDS-PAGE.
<b>Content :</b>	Lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4.
<b>Storage condition :</b>	Lyophilized Flt3 Ligand Rhesus Macaque although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Flt3 Ligand Rhesus Macaque should be stored at 4°C between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.
<b>Amino Acid :</b>	TQDCSFQHSP ISSDFAVKIR ELSDYLLQDY PVTVPSNLQD EELCGALWRL VLAQRWMERL KTVAGSKMQG LLERVNTEIH FVTKCAFQHP PSCLRFVQTN ISRLLQETSE QLVALKPWIT RQNFRCLEL QCQPDSSTLP PPRSPGALEA TALTAPQRP

### Application Note

It is recommended to reconstitute the lyophilized Flt3 Ligand in sterile 18MΩ·cm H<sub>2</sub>O not less than 100µg/ml, which can then be further diluted to other aqueous solutions. The ED50 as calculated by the dose-dependant stimulation of the proliferation of human AML5 cells is less than 1.0ng/ml, corresponding to a specific activity of 1,000,000IU/mg.

