

32-1258: GH 20kDa Recombinant Protein

Alternative Name : GH1,GH,GHN,GH-N,hGH-N,Pituitary growth hormone,Growth hormone 1,Somatotropin.

Description

Source : Escherichia Coli. Growth Hormone 20KDa Pituitary Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 177 amino acids and having a molecular mass of 20322 Dalton. HGH-20kDa is purified by proprietary chromatographic techniques. GH is a member of the somatotropin/prolactin family of hormones which play an important role in growth control. The gene, along with four other related genes, is located at the growth hormone locus on chromosome 17 where they are interspersed in the same transcriptional orientation; an arrangement which is thought to have evolved by a series of gene duplications. The five genes share a remarkably high degree of sequence identity. Alternative splicing generates additional isoforms of each of the five growth hormones, leading to further diversity and potential for specialization. This particular family member is expressed in the pituitary but not in placental tissue as is the case for the other four genes in the growth hormone locus. Mutations in or deletions of the gene lead to growth hormone deficiency and short stature.

Product Info

| | |
|----------------------------|---|
| Amount : | 50 µg |
| Purification : | Greater than 98.0% as determined by:(a) Analysis by SEC-HPLC.(b) Analysis by SDS-PAGE. |
| Content : | HGH-20K was lyophilized from a concentrated (1mg/ml) solution with 0.0045mM NaHCO ₃ previously adjusted pH 10. |
| Storage condition : | Lyophilized Pituitary Growth Hormone-20K although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution and filter sterilization GH-20K can be stored at 4°C for up to 4 weeks. For long term storage and more diluted solutions it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles. |
| Amino Acid : | The sequence of the first five N-terminal amino acids was determined and was found to be Ala-Phe-Pro-Thr-Ile. |

Application Note

It is recommended to reconstitute the lyophilized Growth Hormone Pituitary in 0.4% NaHCO₃ adjusted to pH 10, not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

