

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-1275: raGH Recombinant Protein

Alternative Name: GH1,GH,GHN,GH-N,hGH-N,Pituitary growth hormone,Growth hormone 1,Somatotropin.

Description

Source: Escherichia Coli. Growth Hormone Rabbit Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 190 amino acids and having a molecular mass of 21774 Dalton. GH is purified by proprietary chromatographic techniques. GH is a member of the somatotropin/prolactin family of hormones which play an important role in growth control. The gene, along with four other related genes, is located at the growth hormone locus on chromosome 17 where they are interspersed in the same transcriptional orientation; an arrangement which is thought to have evolved by a series of gene duplications. The five genes share a remarkably high degree of sequence identity. Alternative splicing generates additional isoforms of each of the five growth hormones, leading to further diversity and potential for specialization. This particular family member is expressed in the pituitary but not in placental tissue as is the case for the other four genes in the growth hormone locus. Mutations in or deletions of the gene lead to growth hormone deficiency and short stature.

Product Info

Amount: 20 μg

Purification: Greater than 97.0% as determined by:(a) Analysis by SEC-HPLC.(b) Analysis by SDS-PAGE. **Content:** The protein was lyophilized from a concentrated (1mg/ml) solution with 0.0045mM NaHCO3.

Lyophilized Growth Hormone Rabbit although stable at room temperature for 3 weeks, should

Storage condition : be stored desiccated below -18°C. Upon reconstitution GH Rabbit should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to

add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Application Note

It is recommended to reconstitute the lyophilized Growth Hormone Rabbit in sterile $18M\Omega$ -cm H2O not less than $100\mu g/ml$, which can then be further diluted to other aqueous solutions.

