

32-1302: HGF Sf9 Recombinant Protein

Alternative Name : Scatter Factor (SF),Hepatopoietin (HPTA),HGF,HGFB,F-TCF.

Description

Source : Insect cells. Hepatocyte Growth Factor Human Recombinant produced in Baculovirus is a heterodimer, non-glycosylated, polypeptide chain containing 692 a.a and having a total molecular mass of 78.0 KDa.The HGF is purified by proprietary chromatographic techniques. Hepatocyte Growth Factor (HGF) is a multifunctional growth factor which regulates both cell growth and cell motility. It exerts a strong mitogenic effect on hepatocytes and primary epithelial cells. HGF synergizes with Interleukin-3 and GM-CSF to stimulate colony formation of hematopoietic progenitor cells in vitro and may, therefore, also modulate hematopoiesis.

Product Info

Amount :	10 µg
Purification :	Greater than 95.0% as determined by SDS-PAGE.
Content :	The sterile protein powder (1 mg/ml) is lyophilized from a solution containing 50mM acetic acid.
Storage condition :	Lyophilized Hepatocyte Growth Factor although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution HGF should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
Amino Acid :	QRKRRNTIHEFKKSAKTTLIKIDPALKIKTKKVNTADQCANRCTRNKGLPFTCKAFVFDKARKQCLWFPFNSMSS GVKKEFGHEFDLYENKDYIRNCIIGKGRSYKGTVSITKSGIKQPWSSMIPHEHSYRGKDLQENYCRNPRGEEG GPWCFTSNPEVRYEVCDDIPQCSEVECMTNCGESYRGLMDHTESGKICQRWDHQTPHRHKFLPERYPDKGFDD NYCRNPDGQPRPWICYTLDPHTRWEYCAIKTCADNTMNDTDVPLETTECIQGQGEYRGTVNTIWNIGIPCQR WDSQYPHEHDMTPENFKCKDLRENYCRNPDGSESPWCFTTDPNIRVGYCSQIPNCDMSHGQDCYRGNNGKNY MGNLSQTRSGLTCSMWDKNMEDLHRHIFWEPDASKLNENYCRNPDDAHGPWCYTG NPLIPWDYCPISRCE GDTTPTIVNLDHPVISCATKQLRVVNGIPTRTNIGWMVSLRYRNKHICGGSLIKESWVLTARQC FPSRDLKDYE AWLGIHDVHGRGDEKCKQVLNVSQLVYGPESDLVLMKLARPAVLDDFVSTIDLPNYGCTIPEKTS CSVYVYWG YTGLINYDGLLRVAHLYIMGNEKCSQHHRGKVTLNESEICAGAEEKIGSGPCEGDYGGPLVCEQH KMRMVLGVIV PGRGCAIPNRP GIFVRVAYYAKWIIHKIILTYKVPQS.

Application Note

It is recommended to reconstitute the lyophilized Hepatocyte Growth Factor in 50mM acetic acid to a concentration not less than 100µg/ml. Further dilutions should be made using buffer containing protein. The activity was assayed for scattering activity in the MDCK cell assay. The ED50 for this effect is typically at 1.0-5.0 ng/ml (200,000-1,000,000 IU/mg).

