

## 32-1328: rIFNG Recombinant Protein

**Alternative Name :** Immune Interferon,type II interferon,T cell interferon,MAF,IFNG,IFG,IFI,IFN-gamma.

### Description

Source : Escherichia Coli. Interferon-gamma Rat Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 135 amino acids and having a molecular mass of 15609 Dalton.The IFN-gamma is purified by proprietary chromatographic techniques. IFN-gamma produced by lymphocytes activated by specific antigens or mitogens.IFN-gamma, in addition to having antiviral activity, has important immunoregulatory functions, it is a potent activator of macrophages, and has antiproliferative effects on transformed cells and it can potentiate the antiviral and antitumor effects of the type I interferons.

### Product Info

<b>Amount :</b>	100 µg
<b>Purification :</b>	Greater than 95.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
<b>Content :</b>	Lyophilized from a 0.2µm filtered concentrated (1mg/ml) solution in PBS, pH 7.4.
<b>Storage condition :</b>	Lyophilized Interferon-gamma although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IFN-gamma should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
<b>Amino Acid :</b>	The sequence of the first five N-terminal amino acids was determined and was found to be Met-Gln-Gly-Tyr-Leu.

### Application Note

It is recommended to reconstitute the lyophilized Interferon-gamma in sterile 18MΩ·cm H2O not less than 100µg/ml, which can then be further diluted to other aqueous solutions. The specific activity as determined by the cytopathic affect inhibition assay with murine L929 cells challenged with EMC virus was < 0.1 ng/ml, corresponding to a specific activity of 10,000,000units/mg.

