

## 32-1329: qIFNG Recombinant Protein

**Alternative Name :** Interferon gamma,IFN-gamma,IFNG.

### Description

Source : Escherichia Coli. Recombinant Equine Interferon-gamma produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 143 amino acids and having a molecular mass of 16.7kDa.The IFN-gamma Equine is purified by proprietary chromatographic techniques. IFN-gamma produced by lymphocytes activated by specific antigens or mitogens.IFN-gamma, in addition to having antiviral activity, has important immunoregulatory functions, it is a potent activator of macrophages, and has antiproliferative effects on transformed cells and it can potentiate the antiviral and antitumor effects of the type I interferons.

### Product Info

<b>Amount :</b>	20 µg
<b>Purification :</b>	Greater than 97.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
<b>Content :</b>	Lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4 and 5% trehalose.
<b>Storage condition :</b>	Lyophilized Interferon-gamma although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IFN-gamma should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
<b>Amino Acid :</b>	QA AFFKEIEN LKEYFNASNP DVGDGGLFL DILKNWKEDS DKKIIQSQIV SFYFKLFENL KDNQVIQKSM DTIKEDLFVK FFNSSTSKLE DFQKLIQIPV NDLKVQRKAI SELIKVMNDL SPKANLRKRK RSQNPFRGRR ALQ.

### Application Note

It is recommended to reconstitute the lyophilized Interferon-gamma in sterile 18M-cm H<sub>2</sub>O not less than 100µg/ml, which can then be further diluted to other aqueous solutions. The ED<sub>50</sub> as determined by an anti-viral assay using human HeLa cells infected with encephalomyocarditis (EMC) virus is less than 10.0 ng/ml, corresponding to a specific activity of > 1.0 IU/mg.

