

32-1363: pIL 1 alpha Recombinant Protein

Alternative Name Hematopoietin-1,Lymphocyte-activating factor (LAF),Endogenous Pyrogen (EP),Leukocyte Endogenous Mediator (LEM),Mononuclear Cell Factor (MCF),IL-1 alpha,IL1,IL-1A,IL1F1.

Description

Source : Escherichia Coli. Interleukin-1A Porice Recombinant produced in E.Coli is single, a non-glycosylated, Polypeptide chain containing 158 amino acids and having a molecular mass of 18076 Dalton. The IL-1A is purified by proprietary chromatographic techniques. Interleukin-1 alpha is a proinflammatory cytokine produced by a wide variety of cell types, including macrophages, osteoblasts, monocytes and hepatocytes. Circulating levels of are normally low and only rise after stimulation by agents such as those produced byinflammation, infection or microbial endotoxins. IL-1 alpha possesses a wide variety of biological activities and exerts its effects by binding to specific cell surface receptors.

Product Info

Amount :	10 µg
Purification :	Greater than 95.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
Content :	The protein was lyophilized from a concentrated (1mg/ml) sterile solution containing no additives.
Storage condition :	Lyophilized Interleukin-1 alpha although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL1A should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
Amino Acid :	The sequence of the first five N-terminal amino acids was determined and was found to be Ser-Ala-Thr-Tyr-Ser.

Application Note

It is recommended to reconstitute the lyophilized Interleukin-1 alpha in sterile 18MΩ \square cm H₂O not less than 100 \square µg/ml, which can then be further diluted to other aqueous solutions. The ED50 as determined by the dose-dependant stimulation of D10S cells is < 0.03 ng/ml.

