

## 32-1371: rmlL 1 beta Recombinant Protein

**Alternative Name :** Catabolin,Lymphocyte-activating factor (LAF),Endogenous Pyrogen (EP),Leukocyte Endogenous Mediator (LEM),Mononuclear Cell Factor (MCF),IL1F2,IL-1 beta.

### Description

Source : Escherichia Coli. Interleukin-1 beta Monkey Recombinant produced in E.Coli is a non-glycosylated, Polypeptide chain containing 153 amino acids and having a molecular mass of 17.3 kDa. The IL-1b is purified by proprietary chromatographic techniques. Interleukin-1b is produced by activated macrophages, IL-1B stimulates thymocyte proliferation by inducing il-2 release, b-cell maturation and proliferation, and fibroblast growth factor activity. IL1B proteins are involved in the inflammatory response, being identified as endogenous pyrogens, and are reported to stimulate the release of prostaglandin and collagenase from synovial cells.

### Product Info

**Amount :** 10 µg  
**Purification :** "Greater than 98.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE."  
**Content :** The protein was lyophilized with 1x PBS pH-7.4  
**Storage condition :** Lyophilized IL1B although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL1B should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.  
**Amino Acid :** APVRS LHCTL RDAQLKSLVM SGPYELKALH LQGQDLEQVQV VFSMSFVQGE ESNDKIPVAL GLKAKNLYLS CVLKDDKPTL QLESVDPKNY P KKKMEKRFV FNKIEINNKL EFESAQFPNW YISTSQAENM PVFLGGTRGG QDITDFTMQF VSS.

### Application Note

It is recommended to reconstitute the lyophilized IL1B in sterile 18M $\Omega$ cm H<sub>2</sub>O not less than 100 $\mu$ g/ml, which can then be further diluted to other aqueous solutions. The ED50 as determined by the dose-dependant stimulation of D10.G4.1 mouse helper T cells is typically 3-10pg/mL.

