

32-1385: mL 2 Recombinant Protein

Alternative Name : Interleukin-2,T-cell growth factor (TCGF),Aldesleukin,Lymphokine,IL-2.

Description

Source : Escherichia Coli. Interleukin-2 Mouse Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 148 amino acids and having a molecular mass of 17.2kDa. The IL-2 is purified by proprietary chromatographic techniques. IL2 is a secreted cytokine that is important for the proliferation of T and B lymphocytes. The receptor of this cytokine is a heterotrimeric protein complex whose gamma chain is also shared by interleukin 4 (IL4) and interleukin 7 (IL7). The expression of this gene in mature thymocytes is monoallelic, which represents an unusual regulatory mode for controlling the precise expression of a single gene. The targeted disruption of a similar gene in mice leads to ulcerative colitis-like disease, which suggests an essential role of this gene in the immune response to antigenic stimuli.

Product Info

Amount :	20 µg
Purification :	Greater than 97.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
Content :	Lyophilized from a 0.2µm filtered concentrated solution (1mg/ml) in 20mM PBS, pH 7.4.
Storage condition :	Lyophilized Interleukin-2 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL2 should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
Amino Acid :	PTSSSTSSST AEAQQQQQQ QQQQHLEQL LMDLQELLSR MENYRNLKLP RMLTFKFYLP KQATELKDLO CLEDELGPLR HVLDLTQSKS FQLEDAENFI SNIRVTVVKL KGSDNTFECQ FDDSATVVD FLRRWIAFCQ SIISTSPQ.

Application Note

It is recommended to reconstitute the lyophilized Mouse IL-2 in sterile 18Ml©-cm H2O not less than 100Åµg/ml, which can then be further diluted to other aqueous solutions. The ED50 as determined by the dose-dependant stimulation of murine CTLL-2 cells is less than 0.2ng/ml corresponding to a Specific Activity of 5x106IU/mg.

