

## 32-1387: rIL 2 Recombinant Protein

**Alternative Name :** T-cell growth factor (TCGF),Aldesleukin,Lymphokine,IL-2.

### Description

Source : Escherichia Coli. Interleukin-2 Rat Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 134 amino acids and having a molecular mass of 17kDa. The IL-2 is purified by proprietary chromatographic techniques. IL2 is a secreted cytokine that is important for the proliferation of T and B lymphocytes. The receptor of this cytokine is a heterotrimeric protein complex whose gamma chain is also shared by interleukin 4 (IL4) and interleukin 7 (IL7). The expression of this gene in mature thymocytes is monoallelic, which represents an unusual regulatory mode for controlling the precise expression of a single gene. The targeted disruption of a similar gene in mice leads to ulcerative colitis-like disease, which suggests an essential role of this gene in the immune response to antigenic stimuli.

### Product Info

<b>Amount :</b>	20 µg
<b>Purification :</b>	Greater than 95.0% as determined by SDS-PAGE.
<b>Content :</b>	The protein (1mg/ml) was lyophilized without any additives.
<b>Storage condition :</b>	Lyophilized Interleukin-2 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL2 should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
<b>Amino Acid :</b>	The sequence of the first five N-terminal amino acids was determined and was found to be Met-Ala-Pro-Thr-Ser.

### Application Note

It is recommended to reconstitute the lyophilized Interleukin 2 in sterile 100mM AcOH not less than 100µg/ml, which can then be further diluted to other aqueous solutions. The ED50 as determined by the dose-dependant stimulation of murine CTLL-2 cells is 0.1-0.4 ng/ml corresponding to a specific activity of 2,500,000-10,000,000 units/mg.

