# **w** abeomics

## 32-1392: IL 3 Recombinant Protein

Alternative Name : MCGF (Mast cell growth factor), Multi-CSF, HCGF, P-cell stimulation factor, IL-3, MGC79398, MGC79399.

## Description

Source : Escherichia Coli. Interleukin-3 Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 133 amino acids and having a molecular mass of 15000 Dalton. The IL-3 is purified by proprietary chromatographic techniques. IL3 is a potent growth promoting cytokine. This cytokine is capable of supporting the proliferation of a broad range of hematopoietic cell types. It is involved in a variety of cell activities such as cell growth, differentiation and apoptosis. This cytokine has been shown to also possess neurotrophic activity, and it may be associated with neurologic disorders.

### **Product Info**

Amount : Purification :	10 μg Greater than 98.0% as determined by:(a) Analysis by SEC-HPLC.(b) Analysis by SDS-PAGE.
Content :	Lyophilized from a concentrated (1mg/ml) solution in water containing 0.3 mg/ml of NaHCO3.
Storage condition :	Lyophilized Interleukin-3 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL3 should be stored at 4°C between 2-7 days and for future use below -18°C.Please prevent freeze-thaw cycles.
Amino Acid :	The sequence of the first five N-terminal amino acids was determined and was found to be Ala- Pro-Met-Thr-Gln.

#### **Application Note**

It is recommended to reconstitute the lyophilized Interleukin-3 in sterile  $18M\tilde{A}$   $\tilde{A}$  cm H2O not less than  $100\tilde{A}$   $\tilde{A}\mu g/ml$ , which can then be further diluted to other aqueous solutions. The ED50 as determined by the dose-dependent stimulation of TF-1 cells is < 0.1 ng/ml, corresponding to a Specific Activity of 10,000,000 IU/mg.

