

32-1419: rmlL 6 Recombinant Protein

Alternative Name IFN- β , B cell differentiation factor, BCDF, BSF-2, HPGF, HSF, MGI-2, B-cell stimulatory factor 2, Interferon beta-2, Hybridoma growth factor, CTL differentiation factor, CDF, IL-6, HGF.

Description

Source : Escherichia Coli. IL 6 Rhesus Macaque Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 186 amino acids and having a molecular mass of 21.1kDa. The IL 6 Rhesus Macaque is purified by proprietary chromatographic techniques. IL-6 is a cytokine with a wide variety of biological functions: it plays an essential role in the final differentiation of b-cells into ig-secreting cells, it induces myeloma and plasmacytoma growth, it induces nerve cells differentiation, in hepatocytes it induces acute phase reactants.

Product Info

Amount :	10 μ g
Purification :	Greater than 97.0% as determined by SDS-PAGE and HPLC analyses.
Content :	Lyophilized from a 0.2 μ m filtered concentrated solution in PBS, pH 7.4.
Storage condition :	Lyophilized IL-6 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL-6 should be stored at 4°C between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.
Amino Acid :	MAPVLPGEDS KNVAAPHSQP LTSSERIDKH IRYILDGISA LRKETCNRSN MCESSKEALA ENNLNLPKMA EKDGCFSQSGF NEDTCLVKII TGLLEFEVYL EYLQNRFESE EEQARAVQMS TKVLIQFLQK KAKNLDAITT PEPTTNASLL TKLQANQWL QDMTTHLILR SFKEFLQSNL RALRQM

Application Note

It is recommended to reconstitute the lyophilized IL-6 in sterile 18M-cm H₂O not less than 100 μ g/ml, which can then be further diluted to other aqueous solutions. Fully biologically active when compared to standard. The ED₅₀ determined by a cell proliferation assay using IL-6-dependent murine 7TD1 cells is less than 0.1ng/ml, corresponding to a specific activity of > 1.0 \times 10,000,000 IU/mg.

