∗ abeomics

32-1425: mIL 7 Recombinant Protein

Alternative Name : Lymphopoietin 1 (LP-1), pre-B cell factor, IL-7.

Description

Source : Escherichia Coli. Interleukin-7 Mouse Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 130 amino acids and having a molecular mass of 15.0 kDa. The IL-7 is purified by proprietary chromatographic techniques. Interleukin-7 is a potent lymphoid cell growth factor produced primarily by stromal cells. IL-7 has been shown to support the proliferation and differentiation of pre B- and early T cells as well as displaying a biological effect on cells of NK and myeloid lineages.

Product Info

Amount : Purification : Content :	10 μg Greater than 95.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE. Filtered (0.2μm) and lyophilized from a concentrated (1mg/ml) solution in 1xPBS, pH7.4.
Storage condition :	Lyophilized Interleukin-7 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL7 should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
Amino Acid :	ECHIKDKEGK AYESVLMISI DELDKMTGTD SNCPNNEPNF FRKHVCDDTK EAAFLNRAAR KLKQFLKMNI SEEFNVHLLT VSQGTQTLVN CTSKEEKNVK EQKKNDACFL KRLLREIKTC WNKILKGSI.

Application Note

It is recommended to reconstitute the lyophilized Interleukin 7 in sterile $18M\tilde{A}$ \hat{A} cm H2O not less than $100\tilde{A}$ $\hat{A}\mu g/ml$, which can then be further diluted to other aqueous solutions. The ED50was determined by the dose-dependent stimulation of the proliferation of murine 2E8 cells is less than 0.2 ng/ml, corresponding to a specific activity of >5.0 \tilde{A} \hat{A} $\hat{A}\mu g/ml$, which can the proliferation of murine 2E8 cells is less than 0.2 ng/ml, corresponding to a specific activity of >5.0 \tilde{A} \hat{A}

