

32-1431: IL 10 Recombinant Protein

Alternative Name : B-TCGF,CSIF,TGIF,IL-10,IL10A,MGC126450,MGC126451,Cytokine synthesis inhibitory factor.

Description

Source : Escherichia Coli. Interleukin-10 Human Recombinant produced in E.coli is a single non-glycosylated polypeptide chains containing 161 amino acids each and having a molecular mass of 18.6kDa.The IL-10 is purified by proprietary chromatographic techniques. IL10 is a cytokine produced primarily by monocytes and to a lesser extent by lymphocytes. This cytokine has pleiotropic effects in immunoregulation and inflammation. It down-regulates the expression of Th1 cytokines, MHC class II Ags, and costimulatory molecules on macrophages. It also enhances B cell survival, proliferation, and antibody production. This cytokine can block NF-kappa B activity, and is involved in the regulation of the JAK-STAT signaling pathway. Knockout studies in mice suggested the function of this cytokine as an essential immunoregulator in the intestinal tract.

Product Info

Amount :	10 µg
Purification :	Greater than 97.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
Content :	Lyophilized from a 0.2µm filtered concentrated (1mg/ml) solution in PBS, pH 7.4.
Storage condition :	Lyophilized Interleukin-10 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL10 should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
Amino Acid :	MSPGQGTQSE NSCTHFPGNL PNMLRDLRDA FSRVKTFQMQ KDQLDNLLLK ESLEDFKGY LGCQALSEMI QFYLEEVMPO AENQDPDIKA HVNSLGENLK TLRRLRRCH RFLPCENKSK AVEQVKNAFN KLQEKGIYKA MSEFDIFINY IEAYMTMKIR N.

Application Note

It is recommended to reconstitute the lyophilized Interleukin-10 in sterile 18MΩ·cm H₂O not less than 100µg/ml, which can then be further diluted to other aqueous solutions. The ED50 as determined by the dose-dependent co-stimulation (with murine IL-4) of MC/9 cells was found to be less than 2.0ng/ml, corresponding to a specific activity of 5.0x10⁵ IU/mg.

