

32-1441: IL 12 p40 Baculovirus Recombinant Protein

Alternative Name NKSF2,CTL maturation factor (TCMF),Cytotoxic lymphocyte maturation factor 40 kDa subunit (CLMF : p40),TSF,Edodekin-alpha,IL-12 p40,IL-12B,IL-12 subunit p40,NK cell stimulatory factor chain 2.

Description

Source : Baculovirus IL 12 p40 Human Recombinant produced in Hi-5 cell using Baculovirus is a single polypeptide chain containing 315 amino acids (23-328) and having a molecular mass of 35.8kDa.IL 12 p40 is fused to a 6 amino acid His-tag at C-terminus & purified by proprietary chromatographic techniques. Active IL-12 is a p70 disulphide-linked dimer composed of p35 and p40 subunits. The protein is a pleiotropic cytokine produced primarily by antigen presenting cells and has multiple effects on T lymphocytes and natural killer cells in terms of stimulating cytotoxicity, proliferation, production of other cytokines and Th1 subset differentiation.

Product Info

| | |
|----------------------------|--|
| Amount : | 10 µg |
| Purification : | Greater than 90% as determined by SDS-PAGE. |
| Content : | The IL 12 p40 solution (0.25mg/ml) contains 20mM Tris-HCl buffer (pH 8.0), 100mM NaCl, 2mM DTT, 100mM PMSF and 20% glycerol. |
| Storage condition : | Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Avoid multiple freeze-thaw cycles. |
| Amino Acid : | ADPIWELKGD VYVVELDWYP DAPGEMVVL TCDTPEEDGIT WTL DQSSEVL GSGKTLTIQV KEFGDAGQYT CHKGGEVLSH SLLLLHKKED GIWSTDILKD QKEPKNKTFL RCEAKNYSGR FTCWWLTTIS TDLTFSVKSS RGSSDPQGV TCGAATLSAER VRGDNKEYEY SVECQEDSAC PAAEESLPIE VMVDAVHKLK YENYTSSFFI RDIKPDPPK NLQLKPLKNS RQVEVSWEYP DTWSTPHSYF SLTFCVQVQG KSKREKKDRV FTDKTSATVI CRKNASISVR AQDRYYSSSW SEWASVPCSH HHHHH |

