

32-1473: IL 17F HEK Recombinant Protein(Discontinued)

Alternative Name : Cytokine ML-1,IL-17F,Interleukin-17F precursor,IL17F,ML1,ML-1.

Description

Source : HEK. Interleukin- 17F Human Recombinant produced in HEK cells is a glycosylated homodimer, having a total molecular weight of 38kDa.The IL17F is purified by proprietary chromatographic techniques. IL-17F having an accession number of Q96PD4 is a cytokine that shares sequence similarity with IL17. IL-17F is expressed by activated T cells, and has been shown to stimulate the production of several other cytokines, including IL6, IL8, and CSF2/GM-CSF. IL-17F inhibits the angiogenesis of endothelial cells and induce endothelial cells to produce IL2, TGFB1/TGFB, and monocyte chemoattractant protein-1. IL-17F induces stromal cells to produce proinflammatory and hematopoietic cytokines. Intestinal IL17F gene expression is increased in active CD.IL-17A & IL-17F alleles influence the susceptibility to and pathophysiological features of ulcerative colitis independently. IL-17F and MIF gene polymorphisms are significantly associated with the development of functional dyspepsia. The initiation of IL-17F/IL-17R signaling pathway requires the receptor ubiquitination by TRAF6. IL-17F induces expression of IFN-gamma-inducible protein 10 (IP-10) by activating Raf1-mitogen-activated protein kinase 1/2-extracellular-regulated kinase 1/2-p90 ribosomal S6 kinase-cyclic AMP response element-binding protein signaling pathway.

Product Info

| | |
|----------------------------|---|
| Amount : | 10 µg |
| Purification : | Greater than 95% as observed by SDS-PAGE. |
| Content : | The IL17F was lyophilized from 1mg/ml in 1xPBS. Lyophilized IL-17F although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL17F should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles. |
| Storage condition : | |

Application Note

It is recommended to reconstitute the lyophilized IL-17F in sterile water not less than 100µg/ml, which can then be further diluted to other aqueous solutions. The specific activity was determined by the dose-dependent induction of IL-6 secretion from NHDF Adult fibroblasts, and is typically 100-500ng/ml.

