

32-1489: mIL 20 Recombinant Protein

Alternative Name : IL10D,ZCYTO10,IL-20,MGC96907,Four alpha helix cytokine Zcyto10.

Description

Source : Escherichia Coli. Interleukin-20 Mouse Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 152 amino acids and having a total molecular mass of 17,584 Dalton. The IL-20 is purified by proprietary chromatographic techniques. IL20 is a cytokine structurally related to interleukin 10 (IL10). IL20 cytokine has been shown to transduce its signal through signal transducer and activator of transcription 3 (STAT3) in keratinocytes. A specific receptor for this cytokine is found to be expressed in skin and upregulated dramatically in psoriatic skin, suggesting a role for this protein in epidermal function and psoriasis.

Product Info

Amount :	10 µg
Purification :	Greater than 95.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
Content :	Lyophilized from a concentrated (1mg/ml) solution containing no additives.
Storage condition :	Lyophilized Interleukin-20 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL20 should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
Amino Acid :	The sequence of the first five N-terminal amino acids was determined and was found to be Met-Leu-Lys-Thr-Leu.

Application Note

It is recommended to reconstitute the lyophilized Interleukin-20 in sterile 18MΩ·cm H₂O not less than 100µg/ml, which can then be further diluted to other aqueous solutions. The biological activity was determined by measuring the dose dependent activation of STAT3 in human epidermal keratinocytes (HEK2a cells). Significant STAT3 activation is observed with >100ng/mL of Interleukin-20 recombinant mouse, corresponding to a specific activity of 10,000 units/mg of Interleukin-20 recombinant mouse.

