

32-1508: mL 28B Recombinant Protein

Alternative Name : Interferon lambda-3,IFN-lambda-3,Interleukin-28B,IL-28B,IfnI3,Il28,Il28b.

Description

Source : Escherichia Coli. IL-28B Mouse Recombinant produced in E.coli is a single, non-glycosylated polypeptide chain containing 174 amino acids and having a molecular mass of 19.6kDa.The IL28B is purified by proprietary chromatographic techniques. IL-28B is a cytokine remotely related to type I interferons and the IL-10 family. Interleukin-28B, interleukin 28A (IL28A), and interleukin 29 (IL29) are 3 closely related cytokine genes which form a cytokine gene cluster on a chromosomal region mapped to 19q13. Expression of the cytokines encoded by these 3 genes can be induced by viral infection. All 3 cytokines interact with a heterodimeric class II cytokine receptor which consists of interleukin 10 receptor, beta (IL10RB) and interleukin 28 receptor, alpha (IL28RA).

Product Info

Amount :	20 µg
Purification :	Greater than 97.0% as determined by: (a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
Content :	IL28B protein was lyophilized from a 0.2µm filtered concentrated solution in 1xPBS, pH 7.4. Lyophilized IL-28B although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL28B should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
Storage condition :	
Amino Acid :	DPVPRATRLP VEAKDCHIAQ FKSLSPELQ AFKKAKGAIE KRLEKDMRC SSSLISRAWD LKQLVQQERP KALQAEVALT LKVWENINDS ALTTILGQPL HTLSHIHSQL QTCTQLQATA EPKPPSRRLS RWLHRLQEAQ SKETPGCLEL SVTSNLFQLL LRDLCVAVSG DQCV.

Application Note

It is recommended to reconstitute the lyophilized IL-28B in sterile 18M-cm H2O not less than 100Åµg/ml, which can then be further diluted to other aqueous solutions. The ED50 as determined by an anti-viral assay using human HepG2 cells infected with encephalomyocarditis is less than 30 ng/ml, corresponding to a specific activity of > 3.3 Åµ 104 IU/mg.

