

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-1527: mIL36A Recombinant Protein

Alternative Name: Interleukin 36 alpha,FIL1E,IL1F6,FIL1,IL1(EPSILON),interleukin 1 family member 6 (epsilon),MGC129552,MGC129553.

## **Description**

Source: Escherichia Coli. IL36A Mouse Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 160 amino acids and having a molecular mass of 21.0kDa.The IL36A is purified by proprietary chromatographic techniques. Murine IL-36a belongs to the IL-1 family that includes IL-1b, IL-1a, IL-1a, IL-1a, IL-36ra (IL1F5), IL-36b (IL1F8), IL-36g (IL1F9), IL-37 (IL1F7) and IL-38 (IL-1F10). The IL-1 family members display a 12 b-strand, b-trefoil configuration, and are thought to have ascended from a mutual ancestral gene. Murine IL-36a is a 160 amino acid intracellular and secreted protein which holds no signal sequence, no prosegment and no potential N-linked glycosylation sites. IL-36a is released as a reaction to LPS and the cell ATP-induced activation of the P2X7 receptor. Mouse to human, full length IL-36a/IL-1F6 shares 54% aa sequence homology. IL-36a is mostly found in skin and lymphoid tissues, but also in fetal brain, trachea, stomach and intestine.

## **Product Info**

Amount:  $10 \mu g$ 

**Purification:** Greater than 95.0% as determined by SDS-PAGE.

**Content:** Lyophilized from a 0.2µm filtered concentrated solution in 1xPBS, pH 7.4.

**Storage condition :** Lyophilized IL36A Mouse although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL36A should be stored at 4°C between 2-7 days

and for future use below -18°C.Please prevent freeze-thaw cycles.

Amino Acid: MNKEKELRAA SPSLRHVQDL SSRVWILQNN ILTAVPRKEQ TVPVTITLLP CQYLDTLETN RGDPTYMGVQ

RPMSCLFCTK DGEQPVLQLG EGNIMEMYNK KEPVKASLFY HKKSGTTSTF ESAAFPGWFI AVCSKGSCPL

ILTQELGEIF ITDFEMIVVH

## **Application Note**

It is recommended to reconstitute the lyophilized IL36A in sterile  $18M\tilde{A}\Box\hat{A}\odot$ -cm H2O not less than  $100\tilde{A}\Box\hat{A}\mu g/ml$ , which can then be further diluted to other aqueous solutions. Measured by its binding ability in a functional ELISA to bind recombinant mouse IL-1 Rrp2.

