

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-1529: IL36B Recombinant Protein

Alternative Name Interleukin 36 beta,interleukin 1 family member 8 (eta),Interleukin-1 homolog 2,IL1F8 (Canonical product IL-1F8a),IL-1F8 (FIL1-eta),Interleukin-1 Superfamily e,IL1H2,MGC126880,MGC126882.

Description

Source: Escherichia Coli. IL36B Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 157 amino acids and having a molecular mass of 17.7kDa.The IL36B is purified by proprietary chromatographic techniques. Human IL-36b belongs to the IL-1 family that includes IL-1b, IL-1a, IL-1ra, IL-18, IL-36ra (IL1F5), IL-36b (IL1F8), IL-36g (IL1F9), IL-37 (IL1F7) and IL-38 (IL-1F10). The IL-1 family members display a 12 b-strand, b-trefoil configuration, and are thought to have ascended from a mutual ancestral gene. IL-36 beta is known to be actively secreted. Cells expressing IL-36 beta include resting and activated monocytes and B cells. The receptor for IL-36 beta is a blend of IL-1 Rrp2 and IL-1 RACP. Recombinant IL-36 beta stimulates processes involving NF-kB and MAPK in an IL-1 Rrp2-dependent manner.

Product Info

Amount: $10 \mu g$

Purification: Greater than 95.0% as determined by SDS-PAGE.

Content: Lyophilized from a 0.2µm filtered concentrated solution in 1xPBS, pH 7.4.

Lyophilized IL36B Human although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL36B should be stored at 4°C between 2-7 days

and for future use below -18°C.Please prevent freeze-thaw cycles.

Amino Acid: MNPOREAAPK SYAIRDSROM VWVLSGNSLI AAPLSRSIKP VTLHLIACRD TEFSDKEKGN MVYLGIKGKD

LCLFCAEIQG KPTLQLKEKN IMDLYVEKKA QKPFLFFHNK EGSTSVFQSV SYPGWFIATS TTSGQPIFLT

KERGITNNTN FYLDSVE

Application Note

It is recommended to reconstitute the lyophilized IL36B in sterile 18M-cm H2O not less than $100\tilde{A} \square \hat{A} \mu g/ml$, which can then be further diluted to other aqueous solutions. Measured by its binding ability in a functional ELISA to bind recombinant human IL-1 Rrp2 Fc Chimera.

