

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-1531: mIL36B Recombinant Protein

Alternative Name Interleukin 36 beta,interleukin 1 family member 8 (eta),Interleukin-1 homolog 2,IL1F8 (Canonical product IL-1F8a),IL-1F8 (FIL1-eta),Interleukin-1 Superfamily e,IL1H2,MGC126880,MGC126882.

## **Description**

Source: Escherichia Coli. IL36B Mouse Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 183 amino acids and having a molecular mass of 21.0kDa.The IL36B is purified by proprietary chromatographic techniques. Mouse IL-36b belongs to the IL-1 family that includes IL-1b, IL-1a, IL-1ra, IL-18, IL-36ra (IL1F5), IL-36b (IL1F8), IL-36g (IL1F9), IL-37 (IL1F7) and IL-38 (IL-1F10). The IL-1 family members display a 12 b-strand, b-trefoil configuration, and are thought to have ascended from a mutual ancestral gene. Mouse IL-36b/IL-1F8 is a 183 amino acid protein which holds no signal sequence, no prosegment and no potential N-linked glycosylation site(s). Mouse IL-36b like its human homologue is actively secreted. Mouse IL-36/IL-1F8 shares 61-74% aa homology with human IL-36/IL-1F8 isoform 2 and rat IL-1F8.

## **Product Info**

Amount:  $10 \mu g$ 

**Purification:** Greater than 95.0% as determined by SDS-PAGE.

**Content:** Lyophilized from a 0.2µm filtered concentrated solution in 1xPBS, pH 7.4.

Lyophilized IL36B Mouse although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL36B should be stored at 4°C between 2-7 days

and for future use below -18°C. Please prevent freeze-thaw cycles.

Amino Acid: MMAFPPQSCV HVLPPKSIQM WEPNHNTMHG SSQSPRNYRV HDSQQMVWVL TGNTLTAVPA

SNNVKPVILS LIACRDTEFO DVKKGNLVFL GIKNRNLCFC CVEMEGKPTL OLKEVDIMNL YKERKAOKAF

LFYHGIEGST SVFQSVLYPG WFIATSSIER QTIILTHQRG KLVNTNFYIE SEK

## **Application Note**

It is recommended to reconstitute the lyophilized IL36B in sterile 18M-cm H2O not less than  $100\tilde{A} \square \hat{A} \mu g/ml$ , which can then be further diluted to other aqueous solutions. Measured by its binding ability in a functional ELISA to bind recombinant mouse IL-1 Rrp2.

