

32-1533: IL36G 152 a.a. Recombinant Protein

Alternative Name : Interleukin 36 gamma,IL1F9,interleukin 1 family member 9,Interleukin-1 epsilon,IL-1RP2,IL-1H1,IL1E,interleukin 1-related protein 2,Interleukin-1 homolog 1.

Description

Source : Escherichia Coli. IL36G (152 a.a.) Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 152 amino acids and having a molecular mass of 17.0kDa.The IL36G (152 a.a.) is purified by proprietary chromatographic techniques. IL-36gamma belongs to the IL-1 family which includes IL-1b, IL-1a, IL-1ra, IL-18, IL-36 Ra (IL-1F5), IL-36a (IL-1F6), IL-36b (IL-1F8), IL-37 (IL-1F7) and IL-1F10.). The IL-1 family members display a 12 b-strand, b-trefoil configuration, and are thought to have ascended from a mutual ancestral gene. IL-36g is an 18-22 kDa, 169aa intracellular and secreted protein which holds no signal sequence, no prosegment and no potential N-linked glycosylation sites. Human IL-36g shares 58%- 69% aa sequence homology with mouse, rat, bovine and equine IL-36g, and 23 - 57% aa sequence homology with other family members. The IL-36g receptor is a mixture of IL-1 Rrp2, mostly located in epithelia and keratinocytes, and the extensively expressed IL-1 RAcP. All IL-36 (a, b and g) activate N F-B and MAPK pathways in an IL-1 Rrp2 dependent reaction. Additionally, IL-36g induces production of inflammatory cytokines and chemokines like CXCL8/IL-8.

Product Info

Amount :	10 µg
Purification :	Greater than 95.0% as determined by SDS-PAGE.
Content :	Lyophilized from a 0.2µm filtered concentrated solution in 1xPBS, pH 7.4, containing 5% trehalose.
Storage condition :	Lyophilized IL36g Human although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL36g should be stored at 4°C between 2-7 days and for future use below -18°C.Please prevent freeze-thaw cycles.
Amino Acid :	SMCKPITGTI NDLNQQVWTL QGQNLVAVPR SDSVTPVTVA VITCKYPEAL EQGRGDPIYL GIQNPEMCCLY CEKVGEQPTL QLKEQKIMDL YGQPEPVKPF LFYRAKTGRT STLESVAFPD WFIASSKRDQ PIILTSELGK SYNTAFELNI ND

Application Note

It is recommended to reconstitute the lyophilized IL36g in sterile 18M-cm H2O not less than 100Åµg/ml, which can then be further diluted to other aqueous solutions. Fully biologically active when compared to standard. The ED50 as measured by its ability to induce IL-8 secretion in human preadipocytes is less than 15ng/ml, corresponding to a Specific Activity of 67,000 IU/mg.

