

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-1553: mLeptin Recombinant Protein

Alternative Name: OB Protein, Obesity Protein, OBS, Obesity factor.

Description

Source: Escherichia Coli. Leptin Mouse Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 147 amino acids and having a molecular mass of 16.2 kDa. The Leptin is purified by proprietary chromatographic techniques. A 16-kDa peptide hormone secreted from white adipocytes and implicated in the regulation of food intake and energy balance. Leptin provides the key afferent signal from fat cells in the feedback system that controls body fat stores.

Product Info

Amount: 1mg

Purification: Greater than 95.0% as determined by:(a) Analysis by SEC-HPLC.(b) Analysis by SDS-PAGE. **Content:** The mouse Leptin was lyophilized from a concentrated (1mg/ml) solution containing 0.1% TFA.

Lyophilized Leptin although stable at room temperature for 3 weeks, should be stored

Storage condition : desiccated below -18°C. Upon reconstitution Leptin should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein

(0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Amino Acid: MVPIOKVODD TKTLIKTIVT RINDISHTOS VSAKORVTGL DFIPGLHPIL SLSKMDOTLA VYQQVLTSLP

SONVLOIAND LENLRDLLHL LAFSKSCSLP OTSGLOKPES LDGVLEASLY STEVVALSRL OGSLODILOO

LDVSPEC.

Application Note

It is recommended to reconstitute the lyophilized Leptin in sterile $18M\tilde{A} \square \hat{A} \odot$ -cm H2O not less than $100\tilde{A} \square \hat{A} \mu g/ml$, which can then be further diluted to other aqueous solutions. Biological activity of Mouse Leptin is performed in two different mouse obesity models, ob/ob and NZO. Both strains of mice were treated via intraperitoneal injection once daily at a dose of 5ug Leptin/gram body weight for a period of 14 days. Significant effects on body weight , food consumption, and plasma glucose levels were observed to saline-treated controls.

