

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-1579: LFA 3 Recombinant Protein

Alternative Name: CD58,LFA-3,Ag3,Surface glycoprotein LFA-3.

Description

Source: Chinese Hamster Ovary. Lymphocyte Function-Associated Antigen-3 Fusion Protein Recombinant Human is produced by recombinant DNA technology in a Chinese Hamster Ovary (CHO) mammalian cell expression system. The molecular weight is 91.4 kDa. Recombinant LFA3 is purified by proprietary chromatographic techniques. LFA-3 is ligand of the t-lymphocyte cd2 glycoprotein. This interaction is important in mediating thymocyte interactions with thymic epithelial cells, antigen-independent and dependent interactions of t-lymphocytes with target cells and antigen- presenting cells and the t-lymphocyte rosetting with erythrocytes. In addition, the lfa-3/cd2 interaction may prime response by both the cd2+ and lfa-3+ cells.

Product Info

Amount : 100 μg

Purification: Greater than 98.0% as determined by(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Content: Each mg of CD58 contains 0.8mg sucrose, 0.3mg glycine, 0.25mg sodium citrate dihydrate, and

4µg citric acid monohydrate.

Lyophilized LFA3 Human although stable at room temperature for 3 weeks, should be stored

Storage condition:

desiccated below -18°C. Upon reconstitution Human LFA-3 should be stored at 4°C between 2-7

days and for future use below -18°C. For long term storage it is recommended to add a carrier

protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Application Note

It is recommended to reconstitute the lyophilized LFA-3 Human in sterile $18M\tilde{A}\Box\hat{A}\odot$ -cm H2O not less than $100\tilde{A}\Box\hat{A}\mu g/ml$, which can then be further diluted to other aqueous solutions.

