

**36-1874: Monoclonal Antibody to CD8A (Cytotoxic- & Suppressor T-Cell Marker)(Clone : SPM548)**

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	SPM548
<b>Application :</b>	FACS,IF,IHC
<b>Reactivity :</b>	Human
<b>Gene :</b>	CD8A
<b>Gene ID :</b>	925
<b>Uniprot ID :</b>	P01732
<b>Format :</b>	Purified
<b>Alternative Name :</b>	CD8A,MAL
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	Human CD8 recombinant protein

**Description**


CD8 is a cell surface receptor expressed either as a heterodimer with the CD8 beta chain (CD8 alpha / beta ) or as a homodimer (CD8 alpha / alpha ). A majority of thymocytes and a subpopulation of mature T cells and NK cells express CD8a. CD8 binds to MHC class 1 and through its association with protein tyrosine kinase p56lck plays a role in T cell development and activation of mature T cells. For mature T-cells, CD4 and CD8 are mutually exclusive, so anti-CD8, generally used in conjunction with anti-CD4. It is a useful marker for distinguishing helper/inducer T-lymphocytes, and most peripheral T-cell lymphomas are CD4+/CD8-. Anaplastic large cell lymphoma is usually CD4+ and CD8-, and in T-lymphoblastic lymphoma/leukemia, CD4 and CD8 are often co-expressed. CD8 is also found in littoral cell angioma of the spleen.

**Product Info**

<b>Amount :</b>	100 µg
<b>Purification :</b>	Affinity Chromatography
<b>Content :</b>	100 µg in 500 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
<b>Storage condition :</b>	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

**Application Note**

Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml); ,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),



Formalin-fixed, paraffin-embedded human Tonsil stained with CD8a Monoclonal Antibody (SPM548).