

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-1627: NT 3 Recombinant Protein

Alternative Name: Neurotrophic factor, Nerve growth factor-2, NGF-2, HDNF, NT-3.

Description

Source: Escherichia Coli. Neurotrophin-3 Human Recombinant produced in E.Coli is a non-glycosylated and non-covalently linked homodimer, containing 2x119 amino acid chains, having a total Mw of 27.2 kDa.The NT-3 is purified by proprietary chromatographic techniques. NT3 a member of the neurotrophin family, that controls survival and differentiation of mammalian neurons. This protein is closely related to both nerve growth factor and brain-derived neurotrophic factor. It may be involved in the maintenance of the adult nervous system, and may affect development of neurons in the embryo when it is expressed in human placenta. NTF3-deficient mice generated by gene targeting display severe movement defects of the limbs. The mature peptide of this protein is identical in all mammals examined including human, pig, rat and mouse.

Product Info

Amount: 10 µg

Purification : Greater than 98.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE. **Content :** Lyophilized from a concentrated (1mg/ml) solution in water containing no additives.

Lyophilized NGF2 although stable at room temperature for 3 weeks, should be stored

Storage condition : desiccated below -18°C. Upon reconstitution NGF-2 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein

(0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Amino Acid: YAEHKSHRGE YSVCDSESLW VTDKSSAIDI RGHQVTVLGE IKTGNSPVKQ YFYETRCKEA RPVKNGCRGI

DDKHWNSQCK TSQTYVRALT SENNKLVGWR WIRIDTSCVC ALSRKIGRT.

Application Note

It is recommended to reconstitute the lyophilized Neurotrophin-3 in sterile $18M\tilde{A}\Box\hat{A}\odot$ -cm H2O not less than $100\tilde{A}\Box\hat{A}\mu g/ml$, which can then be further diluted to other aqueous solutions. The ED50 as determined by the dose-dependent proliferation of C6 cells and is $3.6-5.4\tilde{A}\Box\hat{A}\mu g/ml$.

