

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-1647: OSM HEK Recombinant Protein(Discontinued)

Alternative Name: OSM, MGC 20461.

Description

Source: HEK. Oncostatin-M Human Recombinant produced in HEK cells is a glycosylated monomer, having a total molecular weight of 30kDa. The OSM is purified by proprietary chromatographic techniques. Oncostatin M is a member of a cytokine family that includes leukemia-inhibitory factor, granulocyte colony-stimulating factor, and interleukin 6. This gene encodes a growth regulator which inhibits the proliferation of a number of tumor cell lines. It regulates cytokine production, including IL-6, G-CSF and GM-CSF from endothelial cells.

Product Info

Amount: $10 \mu g$

Purification: Greater than 95% as observeed by SDS-PAGE. **Content:** The OSM was lyophilized from 1mg/ml in 1xPBS.

Lyophilized OSM although stable at room temperature for 3 weeks, should be stored desiccated

Storage condition : below -18°C. Upon reconstitution OSM should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or

BSA). Please prevent freeze-thaw cycles.

Application Note

It is recommended to reconstitute the lyophilized OSM in sterile water not less than $100\tilde{A}_{\parallel}\hat{A}\mu g/ml$, which can then be further diluted to other aqueous solutions. The specific activity was determined by the dose-dependent stimulation of the proliferation of human TF-1 cells (human erythroleukemic indicator cell line) and is typically 2-10ng/ml.

