

32-1651: OTOR Recombinant Protein

Alternative Name Otoraplin,Fibrocyte-derived protein,Melanoma inhibitory activity-like protein,OTOR,MIAL,FDP,MIAL1,MGC126737,MGC126739.

Description

Source : Escherichia Coli. Otoraplin Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 111 amino acids and having a molecular mass of 12.7 kDa.The OTOR is purified by proprietary chromatographic techniques. OTOR proteins is also known as fibrocyte-derived protein (Fdp) and Melanoma inhibitory activity-like (MIAL). Otoraplin is a member of the melanoma-inhibiting activity gene family. Otoraplin is a secreted 16 kDa globular protein that is expressed in the inner ear by periotic mesenchyme and developing and mature fibrocytes. OTOR is highly homologous to MIA/cartilage-derived retinoic acid-sensitive protein (CD-RAP), which is a cartilage-specific protein that is also expressed in malignant melanoma cells. The 111 amino acid mature human otoraplin contains 1 SH3 domain (46 - 107 amino acids) and a Tyr at position 50 that is reportedly sulfated. Otoraplin takes part in the initiation of periotic mesenchyme chondrogenesis. Otoraplin is secreted through the Golgi apparatus and plays a role in cartilage development and maintenance. A frequent polymorphism in the translation start codon of OTOR can abolish translation and may be associated with forms of deafness.

Product Info

Amount :	20 µg
Purification :	Greater than 98.0% as determined by(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
Content :	The OTOR protein was lyophilized from a concentrated (1mg/ml) solution containing 20mM PBS pH-7.4 and 130mM NaCl.
Storage condition :	Lyophilized OTOR Recombinant although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution OTOR should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
Amino Acid :	VHGIFMDRLASKKLCADDECVYTISLASAQEDYNAPDCRFINVKKGQIYVYSKLVKENGAGEFWAGSVYGDG QDEMGGVVGYPFRNLVKEQRVYQEATKEVPTTDIDFFCE.

Application Note

It is recommended to reconstitute the lyophilized Otoraplin in sterile 18MΩ·cm H₂O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

