

## 32-1654: PDGF AA HEK Recombinant Protein(Discontinued)

**Alternative Name :** Glioma-derived growth factor,GDGF,Osteosarcoma-derived Growth Factor,ODGF,PDGF-AA,PDGF-1.

### Description

Source : HEK. PDGF-AA Human Recombinant produced in HEK cells is a glycosylated homodimer, having a molecular weight range of 35-45kDa due to glycosylation.The PDGF-AA is purified by proprietary chromatographic techniques. PDGF-AA, PDGF-BB and PDGF-AB, are potent mitogens for a variety of cell types including smooth muscle cells, connective tissue cells, bone and cartilage cells, and some blood cells. The PDGF is stored in platelet alpha-granules and released upon platelet activation. The PDGF is involved in a number of biological processes, including hyperplasia, chemotaxis, embryonic neuron development, and respiratory tubule epithelial cell development. Two distinct signaling receptors used by PDGF have been identified and named PDGFR-alpha and PDGFR-beta. PDGFR-alpha is high-affinity receptor for each of the three PDGF forms. On the other hand, PDGFR-beta interacts with only PDGF-BB and PDGF-AB.

### Product Info

<b>Amount :</b>	10 µg
<b>Purification :</b>	Greater than 95% as observed by SDS-PAGE.
<b>Content :</b>	The PDGF-AA was lyophilized from 1mg/ml in 1xPBS. Lyophilized PDGF-AA although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution PDGF-AA should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
<b>Storage condition :</b>	

### Application Note

It is recommended to reconstitute the lyophilized PDGF-AA in sterile water not less than 100µg/ml, which can then be further diluted to other aqueous solutions. The specific activity was determined by the dose-dependent stimulation of the proliferation of 3T3 cells and is typically 200-1000ng/ml.

