

32-1724: RELM b His Recombinant Protein

Alternative Name : Resistin-like beta,RELm beta,Cysteine-rich secreted protein FIZZ2,Colon and small intestine-specific cysteine-rich protein,Cysteine-rich secreted protein A12-alpha-like 1,Colon carcinoma-related gene protein,RELm-b,XCP2,HXCP2.

Description

Source : Escherichia Coli. RELM-beta Human Recombinant is a His-Tagged Fusion Protein which is 11 kDa protein containing 90 amino acid residues of the RELM-beta human and 12 additional amino acid residues - HisTag . RELM-beta (Resistin-Like Molecule-beta) is a member of a recently identified family of secreted proteins containing a conserved cysteine-rich C-terminus. The RELM family consists of resistin (also called FIZZ3), RELM-alfa (FIZZ1), RELM-beta (FIZZ2) and RELM-gamma. Only resistin and RELM-beta were found in humans whereas all four RELM family members were identified in rodents.RELM-beta appears to be produced as a homodimer exclusively by intestinal goblet cells and can be found in high quantities in stool. Remarkably, stool of germ-free mice displaying sterile intestinal tract does not contain RELM-beta until bacterial colonization takes place after pathogen-free mice entered natural environment. Some, but not all, colon carcinoma cell lines secrete RELM-beta into the cell culture supernatant.The physiological function of RELM-beta is not known. High doses of recombinant RELM-beta showed hyperglycemic effects including lowered glucose disposal and increased hepatic glucose production in mice.

Product Info

Amount : 25 µg
Purification : Greater than 95% as determined by SDS-PAGE.
Content : Filtered and lyophilized from 0.5 mg/ml in 0.05M Acetate buffer pH-4.
Storage condition : Store lyophilized protein at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles. Reconstituted protein can be stored at 4°C for a limited period of time; it does not show any change after two weeks at 4°C.
Amino Acid : MGSTQCSLDS VMDKKIKDVL NSLEYSPSP SKKLSCASVK SQGRPSSCPA GMAVTGCACG YGCGSWDVQLETTCHCQCSV VDWTTCRCCH LTKLRSHHHH HH.

Application Note

Add 0.2 ml of 0.1M Acetate buffer pH-4 and let the lyophilized pellet dissolve completely. For conversion into higher pH value, we recommend intensive dilution by relevant buffer to a concentration of 10⁻⁴µg/ml. In higher concentrations the solubility of this antigen is limited.

