

32-1727: mRELM g Recombinant Protein

Alternative Name : Resistin-like gamma,RELMgamma,RELM-?,RELM-g.

Description

Source : Escherichia Coli. RELM-gamma Mouse Recombinant is a His -Tagged Fusion Protein having a molecular weight of 11 kDa containing 86 amino acid residues of the RELM-gamma Mouse and 16 additional amino acid residues - HisTag . RELM-gamma is a novel member of the resistin-like molecule/found in inflammatory zone (RELM/FIZZ) family in mice and rats. Microarray and real-time RT-PCR experiments revealed a repression of RELMgamma mRNA in nasal respiratory epithelium of cigarette smoke-exposed versus untreated rats. The analysis of the physiological tissue-specific expression revealed highest expression in hematopoietic tissues, suggesting a cytokine-like role for RELM-gamma. RELM-gamma-mRNA is detectable in bone marrow, spleen, and lung as well as in peripheral blood granulocytes. Promyelocytic HL60 cells transfected with a RELM-gamma expression plasmid have an increased proliferation rate compared to mock-transfected cells and display an altered response to retinoic acid-induced granulocytic differentiation. Taken together, these data provide the first experimental evidence that RELM-gamma is a secreted molecule with a restricted expression pattern that may play a role in promyelocytic differentiation.

Product Info

| | |
|----------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Amount : | 25 µg |
| Purification : | Greater than 95% as determined by SDS-PAGE. |
| Content : | Filtered (0.4µm) and lyophilized from 0.5 mg/ml in 0.05M Acetate buffer pH-4. |
| Storage condition : | Store lyophilized protein at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles. Reconstituted protein can be stored at 4°C for a limited period of time; it does not show any change after two weeks at 4°C.The lyophilized protein remains stable until the expiry date when stored at -20°C. |
| Amino Acid : | MRGSHHHHHH GMASHMTLES IVEKKVKELL ANRDDCPSTV TKTFSCSIT ASGRLASCPS GMTVTGCACG YGCGSWDIRD GNTCHCQCST MDWATARCCQ LA. |

Application Note

It is recommended to add 0.1M Acetate buffer pH4 to prepare a working stock solution of approximately 0.5 mg/ml and let the lyophilized pellet dissolve completely. For conversion into higher pH value, we recommend intensive dilution by relevant buffer to a concentration of 10⁻⁶µg/ml. In higher concentrations the solubility of this antigen is limited. Product is not sterile! Please filter the product by an appropriate sterile filter before using it in the cell culture.

