

32-1841: VEGFC Recombinant Protein

Alternative Name : VEGF-C,Vascular endothelial growth factor C,VRP,Flt4 ligand,Flt4-L,Vascular endothelial growth factor-related protein,VEGFC.

Description

Source : Sf9, Insect Cells. VEGF-C Human Recombinant- contains 129 amino acids residues and was fused to a His- tag (6x His) at the C-terminal end. As a result of glycosylation VEGF-C migrates as an 18-24 kDa protein in SDS-PAGE under reducing conditions. VEGF-C, also known as Vascular Endothelial Growth Factor Related Protein (VRP), is a recently discovered VEGF growth factor family member that is most closely related to VEGF-D. Human VEGF-C cDNA encodes a pre-pro-protein of 416 amino acids residues. It is almost identical to the mouse VEGF-C protein. Similar to VEGF-D, VEGF-C has a VEGF homology domain spanning the middle third of the precursor molecule and long N- and C-terminal extensions. In adults, VEGF-C is highly expressed in heart, placenta, ovary and small intestine. Recombinant human VEGF-C, lacking the N- and C-terminal extensions and containing only the middle VEGF homology domain, forms primarily non-covalently linked dimers. This protein is a ligand for both VEGFR-2/KDR and VEGFR-3/FLT-4. Since VEGFR-3 is strongly expressed in lymphatic endothelial cells, it has been postulated that VEGF-C is involved in the regulation of the growth and/or differentiation of lymphatic endothelium. Although recombinant human VEGF-C is also a mitogen for vascular endothelial cells, it is much less potent than VEGF-A.

Product Info

Amount : 10 µg
Purification : Greater than 90.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
Content : Each mg of VEGF-C Human contains 50mg BSA and 1xPBS as buffer.
Storage condition : Lyophilized VEGF-C although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution VEGF-C should be stored at 4°C between 2-7 days and for future use below -18°C.Please prevent freeze-thaw cycles.

Application Note

It is recommended to reconstitute the lyophilized VEGF-C in sterile 18M \square Å©-cm H₂O not less than 100 \square Åµg/ml, which can then be further diluted to other aqueous solutions. Measured by its ability to bind to the VEGFR-3/FLT-4 receptor in the Ba/F3-hVEGFR-3 assay. The ED₅₀ in this assay is about 4ng/ml corresponding to a Specific Activity of 250,000IU/mg.

