

## 32-1870: ENA 78 Recombinant Protein

**Alternative Name :** Small inducible cytokine B5,CXCL5,Epithelial-derived neutrophil-activating protein 78,Neutrophil-activating peptide ENA-78,ENA-78(1-78),chemokine (C-X-C motif) ligand 5,SCYB5.

### Description

Source : Escherichia Coli. Epithelial Neutrophil-Activating Protein 78 Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 74 amino acids and having a molecular mass of 8020 Dalton. The CXCL5 is purified by proprietary chromatographic techniques. Chemokine (C-X-C motif) ligand 5 (CXCL5) is a small cytokine belonging to the CXC chemokine family that is also known as epithelial-derived neutrophil-activating peptide 78 (ENA-78). It is produced following stimulation of cells with the inflammatory cytokines interleukin-1 or tumor necrosis factor-alpha. Expression of CXCL5 has also been observed in eosinophils, and can be inhibited with the type II interferon IFN-. This chemokine stimulates the chemotaxis of neutrophils possesses angiogenic properties. It elicits these effects by interacting with the cell surface chemokine receptor CXCR2. The gene for CXCL5 is encoded on four exons and is located on human chromosome 4 amongst several other CXC chemokine genes. CXCL5 has been implicated in connective tissue remodelling.

### Product Info

<b>Amount :</b>	20 µg
<b>Purification :</b>	Greater than 95.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
<b>Content :</b>	The CXCL5 was lyophilized from a concentrated (1mg/ml) solution in water containing no additives.
<b>Storage condition :</b>	Lyophilized ENA78 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CXCL5 should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
<b>Amino Acid :</b>	The sequence of the first five N-terminal amino acids was determined and was found to be, Ala-Ala -Val-Leu-Arg.

### Application Note

It is recommended to reconstitute the lyophilized ENA-78 in sterile 18MΩ-cm H<sub>2</sub>O not less than 100µg/ml, which can then be further diluted to other aqueous solutions. The biological activity was determined by measuring the dose dependent mobilization of intracellular calcium (calcium flux) with human neutrophils. Significant calcium mobilization is observed with 100ng/mL (corresponding to a Specific Activity of 10,000IU/mg) of recombinant human ENA-78.

