

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-1887: rFractalkine Recombinant Protein

**Alternative Name** Fractalkine,CX3CL1,Neurotactin,CX3C membrane-anchored chemokine,Small inducible cytokine D1,NTN,NTT,CXC3,CXC3C,SCYD1,ABCD-3,C3Xkine.

## **Description**

Source: Escherichia Coli. Fractalkine Rat Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 76 amino acids and having a molecular mass of 8.7kDa.The Fractalkine is purified by proprietary chromatographic techniques. Fractalkine soluble form is chemotactic for t-cells and monocytes, but not for neutrophils. Fractalkine membrane-bound form promotes adhesion of those leukocytes to endothelial cells. Fractalkine regulates leukocyte adhesion and migration processes at the endothelium and binds to CX3CR1. Natural Human Fractalkine is produced as a long protein (373-amino acid) with an extended mucin-like stalk and a chemokine domain on top. The mucin-like stalk permits it to bind to the cell surface. Fractalkine gene is located on human chromosome 16 along with some CC chemokines known as CCL17 and CCL22.

## **Product Info**

**Amount :** 20 μg

**Purification:** Greater than 97.0% as determined by SDS-PAGE.

**Content:** Lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4.

Lyophilized Fractalkine Human although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Fractalkine should be stored at 4°C

between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.

Amino Acid: OHLGMTKCNI TCHKMTSPIP VTLLIHYOLN OESCGKRAII LETROHRHFC ADPKEKWVOD AMKHLDHOTA

ALTRNG

## **Application Note**

It is recommended to reconstitute the lyophilized Fractalkine in sterile  $18M\tilde{A} \square \hat{A} \odot$ -cm H2O not less than  $100\tilde{A} \square \hat{A} \mu g/ml$ , which can then be further diluted to other aqueous solutions. Determined by its ability to chemoattract human monocytes using a concentration range of 5.0-10.0 ng/ml, corresponding to a specific activity of 100,000-200,000 units/mg.

