

32-1888: mFractalkine Recombinant Protein

Alternative Name : Fractalkine,C-X3-C motif chemokine 1,CX3C membrane-anchored chemokine,Neurotactin,Small-inducible cytokine D1,Cx3cl1,Cx3c,Fkn,Scyd1,CX3C,ABCD-3,AB030188,AI848747,D8Bwg0439e.

Description

Source : Escherichia Coli. Fractalkine Mouse Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 76 amino acids and having a molecular mass of 8.7kDa. The CX3CL1 is purified by proprietary chromatographic techniques. Fractalkine soluble form is chemotactic for t-cells and monocytes, but not for neutrophils. Fractalkine membrane-bound form promotes adhesion of those leukocytes to endothelial cells. Fractalkine regulates leukocyte adhesion and migration processes at the endothelium and binds to CX3CR1. Natural Human Fractalkine is produced as a long protein (373-amino acid) with an extended mucin-like stalk and a chemokine domain on top. The mucin-like stalk permits it to bind to the cell surface. Fractalkine gene is located on human chromosome 16 along with some CC chemokines known as CCL17 and CCL22.

Product Info

Amount : 20 µg
Purification : Greater than 97.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
Content : Lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4.
Storage condition : Lyophilized CX3CL1 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CX3CL1 should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
Amino Acid : QHLGMTKCEI MCGKMTSRIP VALLIRYQLN QESCGKRAIV LETTQHRRFC ADPKEKVVQD AMKHLDHQAA ALTKNG.

Application Note

It is recommended to reconstitute the lyophilized CX3CL1 in sterile 18MÅ©-cm H2O not less than 100Åµg/ml, which can then be further diluted to other aqueous solutions. The ED50as determined by a cell proliferation assay using human peripheral blood lymphocytes (PBL) is less than 0.5 Åµg/ml, corresponding to a specific activity of > 2000IU/mg.

