

## 32-1888: mFractalkine Recombinant Protein

**Alternative Name :** Fractalkine,C-X3-C motif chemokine 1,CX3C membrane-anchored chemokine,Neurotactin,Small-inducible cytokine D1,Cx3cl1,Cx3c,Fkn,Scyd1,CX3C,ABCD-3,AB030188,AI848747,D8Bwg0439e.

### Description

Source : Escherichia Coli. Fractalkine Mouse Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 76 amino acids and having a molecular mass of 8.7kDa. The CX3CL1 is purified by proprietary chromatographic techniques. Fractalkine soluble form is chemotactic for t-cells and monocytes, but not for neutrophils. Fractalkine membrane-bound form promotes adhesion of those leukocytes to endothelial cells. Fractalkine regulates leukocyte adhesion and migration processes at the endothelium and binds to CX3CR1. Natural Human Fractalkine is produced as a long protein (373-amino acid) with an extended mucin-like stalk and a chemokine domain on top. The mucin-like stalk permits it to bind to the cell surface. Fractalkine gene is located on human chromosome 16 along with some CC chemokines known as CCL17 and CCL22.

### Product Info

**Amount :** 20 µg  
**Purification :** Greater than 97.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.  
**Content :** Lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4.  
**Storage condition :** Lyophilized CX3CL1 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CX3CL1 should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.  
**Amino Acid :** QHLGMTKCEI MCGKMTSRIP VALLIRYQLN QESCGKRAIV LETTQHRRFC ADPKEKVVQD AMKHLDHQAA ALTKNG.

### Application Note

It is recommended to reconstitute the lyophilized CX3CL1 in sterile 18MÅ©-cm H2O not less than 100Åµg/ml, which can then be further diluted to other aqueous solutions. The ED50as determined by a cell proliferation assay using human peripheral blood lymphocytes (PBL) is less than 0.5 Åµg/ml, corresponding to a specific activity of > 2000IU/mg.

