

32-1968: MIP 3 Alpha Recombinant Protein

Alternative Name :

S Small inducible cytokine A20 precursor,CCL20,Macrophage inflammatory protein 3 alpha,MIP-3-alpha,Liver and activation- regulated chemokine,CC chemokine LARC,Beta chemokine exodus-1,CKb4,LARC,ST38,MIP3A,MIP-3a,SCYA20.

Description

Source : Escherichia Coli. MIP-3 Alpha Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 70 amino acids and having a molecular mass of 8 kDa. The MIP-3a is purified by proprietary chromatographic techniques. CCL-20 is a chemotactic factor that draws lymphocytes & neutrophils, rather than monocytes. MIP-3 alpha inhibits proliferation of myeloid progenitors in colony formation assays. MIP3A plays a role in the formation and function of the mucosal lymphoid tissues by attracting lymphocytes and dendritic cells towards epithelial cells. C-terminal processed forms have been shown to be equally chemotactically active for leukocytes. CCL-20 holds antibacterial activity e.coli atcc 25922 and s.aureus atcc 29213. CCL-20 gene transcription is activated by H. pylori, which activates NF-kappaB through intracellular signal pathway which involves IkappaB kinase and NF-kappaB-inducing kinase. MIP-3 alpha is involved in chemokine-mediated lymphocyte trafficking during gastric inflammation in Helicobacter infection. CCL-20 expression is involved in the recruitment of CD45R0-positive T cell subsets into the intestinal lamina propria. MIP-3A is in charge of the advancement of pulpal inflammation through the recruitment of C-C motif Receptor 6-expressing lymphocytes Vaginal epithelial cells respond to factors present in semen by secreting MIP-3 alpha, which increases langerhans cells recruitment during HIV transmission.

Product Info

| | |
|----------------------------|--|
| Amount : | 20 µg |
| Purification : | Greater than 97.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE. |
| Content : | Lyophilized from a 0.2µm filtered concentrated solution in 20mM PB, pH 7.4, 150mM NaCl. |
| Storage condition : | Lyophilized MIP-3a although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CCL20 should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles. |
| Amino Acid : | ASNFDCCCLGY TDRILHPKFI VGFTRQLANE GCDINAIIFH TKKKLSVCAN PKQTWVKYIV RLLSKKVKNM. |

Application Note

It is recommended to reconstitute the lyophilized MIP-3 Alpha in sterile 18MΩ-cm H₂O not less than 100µg/ml, which can then be further diluted to other aqueous solutions. Determined by its ability to chemoattract human T lymphocytes using a concentration range of 10.0 -50.0 ng/ml, corresponding to a specific activity of 20,000-100,000units/mg.

