

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-1979: PF 4 Recombinant Protein

Alternative Name: CXCL4,PF-4,PF4,Iroplact,Oncostatin-A,SCYB4,MGC138298.

Description

Source: Escherichia Coli. CXCL4 Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 70 amino acids and having a molecular mass of 7.8 kDa. Platelet factor-4 is a 70-amino acid protein that is released from the alpha-granules of activated platelets and binds with high affinity to heparin. Its major physiologic role appears to be neutralization of heparin-like molecules on the endothelial surface of blood vessels, thereby inhibiting local antithrombin III activity and promoting coagulation. As a strong chemoattractant for neutrophils and fibroblasts, PF4 probably has a role in inflammation and wound repair. Oncostatin-A is a member of the CXC chemokine family. Human PF4 is used for the proof of heparin-induced thrombocytopenia. Furthermore it is used as an inhibitor in the angiogenesis during tumor therapy.

Product Info

Amount : 20 μg

Purification : Greater than 95.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Content: The CXCL4 protein was lyophilized after extensive dialysis against 50mM Tris-HCl pH 8.0 and

150mM NaCl buffer.

Lyophilized CXCL4 although stable at room temperature for 3 weeks, should be stored

Storage condition:

desiccated below -18°C. Upon reconstitution CXCL4 should be stored at 4°C between 2-7 days

and for future use below -18°C.For long term storage it is recommended to add a carrier protein

(0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Amino Acid: The sequence of the first four N-terminal amino acids was determined and was found to be Glu-

Ala-Glu-Glu-Asp.

Application Note

It is recommended to reconstitute the lyophilized CXCL4 in sterile 18MÃ \square Â \square -cm H2O not less than 100Ã \square Â μ g/ml, which can then be further diluted to other aqueous solutions. The ED50 of CXCL4 as determined by its ability to inhibit human FGF basic dependent proliferation of NR6R3T3 mouse fibroblasts was found to be 5-15 Ã \square Â μ g/ml.

