

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

36-1952: Monoclonal Antibody to CD59 / Complement Regulatory Protein / Protectin(Clone : BRA-10G)

Clonality: Monoclonal Clone Name: BRA-10G

Application: Functional Assay, FACS, IF

Reactivity: Human
Gene: CD59
Gene ID: 966
Uniprot ID: P13987
Format: Purified

Alternative Name: CD59,MIC11,MIN1,MIN2,MIN3,MSK21

Isotype: Mouse IgG1, kappa **Immunogen Information:** Human K562 tumor cells

Description

Reacts with human CD59, a 20kDa glycosyl phosphatidyl-inositol (GPI)-anchored cell surface protein. CD59 regulates complement-mediated cell lysis, and it is involved in lymphocyte signal transduction. This protein is a potent inhibitor of the complement membrane attack complex, whereby it binds complement C8 and/or C9 during the assembly of this complex, thereby inhibiting the incorporation of multiple copies of C9 into the complex, which is necessary for osmolytic pore formation. CD59 is widely distributed on cells in all tissues. It inhibits formation of MAC, thus protecting cells from complement-mediated lysis. The expression of CD59 on erythrocytes is important for their survival. Genetic defects in GPI-anchor attachment, that cause a reduction or loss of CD59 and CD55 on erythrocytes produce the symptoms of the disease paroxysmal hemoglobinuria (PNH). This MAb is useful for study on GPI-anchored proteins, PNH and CD59 functions.

Product Info

Amount: 100 μg

Purification: Affinity Chromatography

Content: 100 μg in 500 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly

toxic.

Storage condition : Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Functional Studies (Order Ab without Azide); Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml);