

36-1957: Monoclonal Antibody to CD63 (Late Endosomes Marker)(Clone : NKI/C3)-PE

Clonality :	Monoclonal
Clone Name :	NKI/C3
Application :	FACS,IF
Reactivity :	Human, Mouse
Conjugate :	PE
Gene :	CD63
Gene ID :	967
Uniprot ID :	P08962
Format :	Purified
Alternative Name :	CD63,MLA1,TSPAN30
Isotype :	Mouse IgG1, kappa
Immunogen Information :	Smooth plasma membrane fraction of MeWo cells

Description


This MAb recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of MAb LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multi-molecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression.

Product Info

Amount :	0.5 ml
Purification :	Affinity Chromatography
Content :	PE conjugated CD63 Prepared in 10mM PBS with 0.05% BSA and 0.05% azide.
Storage condition :	Store the antibody at 4°C; stable for 6 months.

Application Note

Flow Cytometry (5ul per test per one million cells or 5ul per 100ul of whole blood);,Immunofluorescence (1:50-1:100 for 30 minutes at RT);,



Flow Cytometry of human CD63 on MCF-7 Cells. Black: Cells alone; Green: Isotype Control; Red: PE-labeled CD63 Monoclonal Antibody (NK1/C3).