

36-11032: Monoclonal Antibody to Clone : IPO-38 (Proliferation Marker)(Clone : IPO-38)

Clonality :	Monoclonal
Clone Name :	IPO-38
Application :	IHC
Reactivity :	Human, Mouse, Rat
Format :	Purified
Isotype :	Mouse IgM, kappa
Immunogen Information :	Spleen cells of a patient with hairy cell leukemia

Description


Recognizes a protein of 14-16kDa, which is a novel nuclear antigen of proliferating cells. IPO-38 antigen is present in the nuclei of proliferating cells such as Hodgkin s disease and non-Hodgkin s lymphomas, different forms of leukemias, breast and colorectal carcinomas, and PHA-stimulated lymphocytes. It is not expressed in the cells of non-stimulated lymphocytes and granulocytes. IPO-38 may be a useful marker of cell proliferation during monitoring of tumor progression.

Product Info


Amount :	100 µg
Purification :	Affinity Chromatography
Content :	100 µg in 500 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
Storage condition :	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

Application Note

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);



Formalin-fixed, paraffin-embedded human Tonsil stained with IPO-38 Monoclonal Antibody.



Formalin-fixed, paraffin-embedded human Tonsil stained with IPO-38 Monoclonal Antibody.