

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

36-11073: Monoclonal Antibody to Melanoma Marker (MART-1 + gp100 + Tyrosinase)(A103 + HMB45 + T311)

Clonality: Monoclonal

Clone Name: A103 + HMB45 + T311

Application: WB,IHC
Reactivity: Human
Gene: MLANA
Gene ID: 2315
Uniprot ID: Q16655
Format: Purified
Alternative Name: MLANA.MART1

Isotype: Mouse IgG1, kappa + Mouse IgG2a, kappa + Mouse IgG1, kappa

Immunogen Information: Recombinant hMART-1 protein (A103); Recombinant tyrosinase protein (T311); Extract of

pigmented melanoma metastases from lymph nodes (HMB45)

Description

This antibody cocktail recognizes three melanoma-specific proteins, which include MART-1, Tyrosinase and gp100. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Tyrosinase is one of the targets for cytotoxic T-cell recognition in melanoma patients. The function of gp100 is not known but it is reported to be a useful marker for melanocytes and melanomas. This cocktail of three markers is designed for extremely sensitive labeling of formalin-fixed, paraffin-embedded melanomas and other tumors showing melanocytic differentiation.

Product Info

Amount: $100 \mu g$

Purification: Affinity Chromatography

Content: 100 μg in 500 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly


toxic.

Storage condition : Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Western Blot (2-4ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);



Formalin-fixed, paraffin-embedded human Melanoma stained with Melanoma Marker Monoclonal Antibody (A103 + T311 + HMB45).