

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

10-4125: Monoclonal antibody to Human CD25 (Clone: 7G7B6)

Clonality: Monoclonal **Clone Name:** 7G7B6 FACS Application: Reactivity: Human Gene: IL2RA Gene ID: 3559 **Uniprot ID:** P01589 **Purified** Format: **Alternative Name:** IL2RA,CD25

Isotype: Mouse IgG2a Kappa

Immunogen Information: Recombinant human CD25 protein was used as an imunogen for this antidody.

Description

CD25 is the alpha chain of the IL-2 receptor. It is a type I transmembrane protein present on activated T cells, activated B cells, some thymocytes, myeloid precursors, and oligodendrocytes that associates with CD122 to form a heterodimer that can act as a high-affinity receptor for IL-2. Though CD25 has been shown to influence effector function of lymphocytes, it is thought to play a greater role in immune tolerance in mice. Constitutive expression of CD25 on Treg cells is thought to be crucial to their survival and maintenance of immune homeostasis. Expression of CD25 and its role in immunology may be species dependent, since CD25 appears to play a larger role in T-cell effector responses in humans compared to mice, and may be somewhat dispensable for the maintenance of Treg cells as seen in patients treated with CD25-blocking antibodies.

Product Info

Amount : $25 \mu g / 100 \mu g$

Purification: Protein G Chromatography

Content: 25 μg in 50 μl/100 μg in 200 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium

azide is highly toxic.

Storage condition : Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

FACS Analysis: 0.5-1 µg/10^6 cells



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig:1- Cell surface flow analysis of hCD25 in human PBMC using 0.5 μ g/10^6 cells. Green represents isotype control (ABEOMICS); red represents anti-hCD25 antibody (10-4125). Goat anti-mouse PE conjugated secondary antibody (ABEOMICS) was used.