

## 10-3012: Monoclonal antibody to MyD88 (Clone: ABM2H20)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	ABM2H20
<b>Application :</b>	FACS, WB
<b>Reactivity :</b>	Mouse, Human
<b>Gene :</b>	MYD88
<b>Gene ID :</b>	4615
<b>Uniprot ID :</b>	Q99836
<b>Format :</b>	Purified
<b>Alternative Name :</b>	Myeloid differentiation primary response protein MyD88
<b>Isotype :</b>	Mouse IgG1 Kappa
<b>Immunogen Information :</b>	A partial length recombinant protein of human MyD88 (amino acids 13-221) was used as an immunogen for this antibody.

### Description

MyD88 (Myeloid differentiation factor) is an essential adaptor molecule in all TLR (Toll-like receptor) signaling pathways except TLR3. MyD88 is composed of an N-terminal (Death Domain) and a highly conserved C-terminal TIR (Toll/interleukin-1 Receptor) domain. It is found to stimulate IL-1R/IL18R-mediated signaling. MyD88-dependent signaling is also important in the regulation of innate as well as acquired immunity, in particular, T-cell responses, to various microbial pathogens. After activation of TLRs, MyD88 is phosphorylated and subsequently recruits IRAKs (IL-1R Associated Kinases) and other downstream proteins such as TRAF6, finally resulting in activation of the NF-kappaB (nuclear factor kappa B) pathway.

### Product Info

<b>Amount :</b>	25 µg / 100 µg
<b>Purification :</b>	Protein G Chromatography
<b>Content :</b>	25 µg in 50 µl/100 µg in 200 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
<b>Storage condition :</b>	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

### Application Note

Western blot analysis: 0.1-2 µg/ml; FACS Analysis: 0.5-1 µg/ml




Fig. 1: Intracellular flow analysis of MyD88 in Jurkat using 0.5  $\mu\text{g}/10^6$  cells of MyD88 antibody (Clone: ABM2H20). Green represents isotype control; red represents anti-MyD88 antibody. Goat anti-mouse PE conjugate was used as secondary antibody.




Fig. 2: Western blot analysis of MyD88. Anti-MyD88 antibody (Clone: ABM2H20) was tested at 0.1  $\mu\text{g}/\text{ml}$  on h Kidney lysate.




Figure:3- Western blot analysis of MyD88. Anti-MyD88 antibody (Clone: ABM2H20) was tested at 1  $\mu\text{g}/\text{ml}$  on (1) Raw and (2) EL-4 lysates.