

## 14-100ACL: NF-kB LEEporter™ Luciferase Reporter-RAW264.7 Cell Line

**Application :** Functional Assay

### Description

The NF-kB LEEporter™ Luciferase Reporter cell line is a stably transfected RAW 264.7 cell line which expresses Renilla luciferase reporter gene under the transcriptional control of the NF-kB response element. NF-kB is a key transcription factor that is involved in immune and inflammatory responses, developmental processes, cellular growth and apoptosis.

### Product Info

**Amount :** 1 vial  
**Content :** Each vial contains  $2 \sim 3 \times 10^6$  cells in 1 ml of 90% FBS + 10% DMSO.  
**Storage condition :** Immediately upon receipt, store in liquid nitrogen.

### Application Note

#### Application:

- Monitor the NF-kB signaling pathway.
- Screen for activators or inhibitors of the NF-kB signaling pathway.

#### Culture conditions:

Cells should be grown at 37°C with 5% CO<sub>2</sub> using DMEM medium (w/ L-Glutamine, 4.5g/L Glucose and Sodium Pyruvate) supplemented with 10% heat-inactivated FBS and 1% Pen/Strep, plus 3 µg/ml of Puromycin (Note: Puromycin can be omitted during the reporter cell assays).

It is recommended to quickly thaw the frozen cells upon receipt or from liquid nitrogen in a 37°C water-bath, transfer to a tube containing 10 ml of growth medium without Puromycin, spin down cells, resuspend cells in pre-warmed growth medium without Puromycin, transfer resuspended cells to T25 flask and culture in 37°C-CO<sub>2</sub> incubator.

Leave the T25 flask in the incubator for 1~2 days without disturbing or changing the medium until cells completely recover viability and become adherent. Once cells are over 90% adherent, remove growth medium and passage the cells through trypsinization and centrifugation. At first passage, switch to growth medium containing Puromycin. Cells should be split before they reach complete confluence. **Note: RAW264.7 cells may not be detached well by trypsinization only. So you may need to use a cell scraper to harvest the trypsinized cells.**

To passage the cells, detach cells from culture vessel with Trypsin/EDTA, add complete growth medium and transfer to a tube, spin down cells, resuspend cells and seed appropriate aliquots of cells suspension into new culture vessels. Subcultivation ration = 1:10 to 1:20 weekly. To achieve satisfactory results, cells should not be passaged over 16 times.

#### Functional validation:

##### A. Response of NF-kB LEEporter™ – RAW264.7 cells to lipopolysaccharide (LPS)

1. Plate NF-kB LEEporter™ – RAW264.7 cells into a white solid-bottom 96-well microplate in 100 µl of growth medium

at  $1 \times 10^5$  cells/well and incubate cells at 37°C in a CO<sub>2</sub> incubator for 4-6 hours.

2. Stimulate cells with different concentrations of LPS and incubate cells at 37°C in a CO<sub>2</sub> incubator for 16 hours. ,

3. Equilibrate the plate to room temperature for 10 minutes.

4. Add 50 µl of luciferase assay reagent (Abeomics, Cat #17-1101; Refer to the reagent datasheet for the detailed luciferase assay protocol) per well.

5. Read the plate in 1-5 minutes to measure luminescence using a microplate luminometer.

#### LIMITED USE RESTRICTIONS:

**THIS PRODUCT IS SOLELY FOR IN VITRO RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.**

**By use of this product, user agrees to be bound by the terms of this limited use statement.**

**This product is solely for Internal Research Purposes and not for Commercial Purposes. Commercial Purposes include, but are not limited to (1) use of the cell line in manufacturing; (2) use of the cell line to provide a service, information or data; (3) use of the cell line for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the cell line whether or not such cell lines are resold for use in research. The buyer cannot sell, give or otherwise transfer this product to a third party.**

**Commercial License Agreement is available for non-research/ non-inhouse/ commercial use if applicable. Please contact Abeomics ([info@abeomics.com](mailto:info@abeomics.com)).**




fig-1: Induction of NF-kB activity by LPS (Cat. No. 15-1013) in NF-kB LEEporter™ - RAW264.7 cells.